

**Classification and synonymy of the crane-flies
described by Enrico Brunetti (Diptera:
Families Ptychopteridae, Trichoceridae
and Tipulidae**

By

CHARLES P. ALEXANDER

Reprinted from:

Records of the Indian Museum, Vol. 59 (Parts 1 & 2)
[for March-June, 1961]

Pages 19-34

PRINTED IN INDIA, BY THE MANAGER, GOVERNMENT OF INDIA PRESS, CALCUTA,
AND PUBLISHED BY THE MANAGER OF PUBLICATIONS, CIVIL LINES, DELHI, 1963.

PUBLISHED: 25th March, 1963

CLASSIFICATION AND SYNONYMY OF THE CRANE-FLIES
DESCRIBED BY ENRICO BRUNETTI (DIPTERA: FAMILIES
PTYCHOPTERIDAE, TRICHOCERIDAE AND TIPULIDAE)

By

CHARLES P. ALEXANDER

Amherst, Massachusetts

CONTENTS

I—Introduction	19
II—The Indian species of crane-flies described by Brunetti	20
Family Ptychopteridae	20
Family Trichocerida	21
Family Tipulidae	21
III—The extra-limital Tipulidae described by Brunetti	33
IV—Synopsis	34

I—INTRODUCTION

Between 1911 and 1924, the late Enrico Brunetti described as new approximately 240 species of crane-flies, the great majority from the area comprised within the limits of former British India and Burma. Due in part to circumstances beyond his control, as discussed in papers by Alexander (1942) and Edwards (1924)**, Brunetti's arrangement of many of his species to genera was questionable, with the result that subsequent taxonomic work on these flies from this faunal area proved difficult and uncertain. Edwards (1924) was able to examine certain of the Brunetti types that were brought to London by the describer while some others have been seen by the writer, having been loaned me for study through the kind and appreciated interest of the authorities of the Indian Museum. It is believed that the present distribution of these species, with an indication of synonymy as it appears certain at the present time, may prove helpful to later workers on the crane-flies of this particularly interesting region.

*Contribution No. 1234 from the Department of Entomology, University of Massachusetts, Amherst, Massachusetts, U.S.A.

**Dates in parenthesis refer in most instances to papers cited in the short Bibliography that follows this introductory statement.

Almost all of the species concerned are Indian and these are treated separately from the few further forms that were described from elsewhere in the Oriental and Australasian regions. In virtually all cases, the locality given represents the type station for the species in question.

BIBLIOGRAPHY

- ALEXANDER, CHARLES P. 1913. The Tipulidae in Brunetti's "Fauna of British India ; Diptera Nematocera".—*Insec. Inscit. Menst.*, Washington, 1, pp. 118-120.
- ALEXANDER, CHARLES P. 1942. The Oriental Tipulidae in the collection of the Indian Museum. Part II.—*Rec. Indian Mus.*, Delhi, 44, pp. 29-72, 21 figs.
- BRUNETTI, E. 1911. Revision of the Oriental Tipulidae with descriptions of new species.—*Rec. Indian Mus.*, Calcutta, 6, pp. 231-314.
- BRUNETTI, E. 1912. The Fauna of British India including Ceylon and Burma. Dipater, Nematocera (excluding Chironomidae and Culicidae), pp. xxx+581, 44 figs., 8 pls. (Tipuloidea, pp. 265-548, 562-572).
- BRUNETTI, E. 1913a. Zoological results of the Abor expedition, 1911-12. XI. Diptera.—*Rec. Indian Mus.*, Calcutta, 8, pp. 149-190, 1 pl.
- BRUNETTI, E. 1913b. New and interesting Diptera from the eastern Himalayas.—*Rec. Indian Mus.*, Calcutta, 9, pp. 255-277, 1 pl.
- BRUNETTI, E. 1918. Revision of the Oriental Tipulidae with descriptions of new species. Part II.—*Rec. Indian Mus.*, Calcutta, 15, pp. 255-340, 7 figs., 2 pls.
- BRUNETTI, E. 1920. Catalogue of Oriental and South Asiatic Nemocera.—*Rec. Indian Mus.*, Calcutta, 17, pp. 1-300 (Includes alphabetical list of localities, pp. 2-8).
- BRUNETTI, E. 1924. Diptera of the Siju cave, Garo Hills, Assam. I. Tipulidae, Tabanidae, Anthomyidae, Acalyptrate Muscidae and Phoridae.—*Rec. Indian Mus.*, Calcutta, 26, pp. 99-106.
- EDWARDS, FRED W. 1924. Notes on the types of Diptera Nematocera (Mycetophilidae and Tipulidae) described by Mr. E. Brunetti.—*Rec. Indian Mus.*, Calcutta, 26, pp. 291-307.

II—THE INDIAN SPECIES OF CRANE-FLIES DESCRIBED BY BRUNETTI

PTYCHOPTERIDAE

- | | | |
|---|-----------|-------------------------------|
| * <i>Ptychoptera annandalei</i> Brun., 1918 : 256 | | Burma : Southern Shan States. |
| <i>P. distincta</i> Brun., 1911 : 232 | | E. Himalayas : Darjiling. |
| * <i>P. tibialis</i> Brun., 1911 : 233 (synonym <i>P. atritarsis</i> Brun., 1911 : 234) | | E. Himalays : Darjiling. |

*Indicates species of which type material is present in the collection of the Zoological Survey of India, Calcutta (A.P. Kapur).

TRICHOCERIDAE

- **Paracladura elegans* Brun., 1911 : 288 . . . E. Himalayas : Darjiling.
 **P. gracilis* Brun., 1911 : 287 (synonym *P. flava* Brun., 1912 : 512, as *Trichocera*) . . . E. Himalayas : Darjiling.
 **Trichocera montana* Brun., 1912 : 513 . . . W. Himalayas : Garwal District.
 **T. punctipennis* Brun., 1912 : 511 . . . W. Himalayas : Simla.

TIPULIDAE

TIPULINAE

- Ctenacroscelis carmichaeli* (Brun., 1913 b : 257, as *Tipula*) . . . E. Himalayas : Darjiling.
 **C. dives* (Brun., 1912 : 307, as *Tipula*) . . . E. Himalayas : Kurseong.
 **C. flavoides* (Brun., 1918 : 267, as *Tipula*) . . . Assam : Garo Hills.
C. flavus (Brun., 1911 : 252, as *Tipula*) . . . E. Himalayas : Sikkim.
 **C. fulvolateralis* (Brun., 1911 : 249, as *Tipula*) (synonym *C. sikkimensis* Enderlein, 1912 : 1) . . . W. Himalayas : Kumaon.
C. fumipennis (Brun., 1911 : 250, as *Tipula*) (pre-occupied—renamed *cerbereanus* Alexander, 1942 : 29) . . . E. Himalayas : Darjiling.
 **C. majesticus* (Brun., 1911 : 248, as *Tipula*) . . . E. Himalayas : Darjiling.
C. ochripes (Brun., 1911 : 260, as *Tipula*) . . . Ceylon.
 **C. ornatithorax* (Brun., 1911 : 258, as *Tipula*) . . . E. Himalayas : Darjiling.
 **C. serricornis* (Brun., 1912 : 309, as *Tipula*) (pre-occupied—renamed *cressida* Alexander, 1953) . . . W. Himalayas : Kumaon.
 **Nephrotoma concolorithorax* (Brun., 1912 : 346, as *Pachyrhina*) . . . Assam : Khasi Hills.
 **N. consimilis* (Brun., 1911 : 266, as *Pachyrhina*) . . . E. Himalayas : Darjiling.
 **N. dorsopunctata* (Brun., 1911 : 265, as *Pachyrhina*) . . . Bihar : Purneah District.
N. fuscoflava (Brun., 1918 : 275, as *Pachyrhina*) . . . Pakistan : Punjab.
 **N. gamma* (Brun., 1912 : 347, as *Pachyrhina*) . . . Assam-Bhutan Border.
 **N. hypocrites* (Brun., 1918 : 276, as *Pachyrhina*) (synonym—*N. nigroapicalis* Brun., 1918 : 335, as *Eriocera*) . . . E. Himalayas : Darjiling.
N. parvinotata (Brun., 1918 : 276, as *Pachyrhina*) . . . Pakistan : Peshawar District.
 **N. pleurinotata* (Brun., 1912 : 343, as *Pachyrhina*) . . . Ceylon.
 **N. punctieornis* (Brun., 1912 : 343, as *Pachyrhina*) . . . E. Himalayas : Darjiling.
 **N. serricornis* (Brun., 1912 : 341, as *Pachyrhina*) . . . E. Himalayas : Darjiling.

- **Pachyrhina demarcata* Brun., 1912 : 344 (pre-occupied—renamed *Tipula (Schummelia) sessilis* Edwards) E. Himalayas Darjiling.
- Ctenophora (Tanyptera) indica* (Brun., 1918 : 257, as *Xiphura*) E. Himalayas : Darjiling.
- **C. (Pselliophora) aurantia* (Brun., 1918 : 260, as *Pselliophora*) Assam Garo Hills..
- **C. (P.) approximata* (Brun., 1918 : 259, as *Pselliophora*) S. India North Kanara District.
- **C. (P.) divisa* (Brun., 1911 : 242, as *Pselliophora*) "East Indies" E. Himalayas.
- **C. (P.) flavofasciata* (Brun., 1918 : 259, as *Pselliophora*) Assam Garo Hills.
- **C. (P.) immaculipennis* (Brun., 1911 : 243, as *Pselliophora*) Assam : Sylhet.
- C. (P.) laeta trilineata* (Brun., 1911 : 240, as *Pselliophora*) S. India Nilgiri Hills.
- C. (P.) latifascipennis* (Brun., 1918 : 260, as *Pselliophora*) S. India Coorg.
- Prionota (Plocimas) serraticornis* (Brun., 1911 : 242, as *Pselliophora*) (synonym *P. (P.) elongata* Edwards, 1913 : 202, as *Pselliophora*) Ceylon.
- **Tipu a (Brithura) gravelvi* Brun., 1918 : 264 (synonym of *T. (B.) crassa* Edwards, 1916) E. Himalayas : Darjiling.
- T. (B.) imperfecta* Brun., 1913 : 260 E. Himalayas Darjiling.
- **T. (Bellardina) griseipennis* Brun., 1912 : 321 W. Himalayas Garhwal District.
- **T. (B.) splendens* Brun., 1912 : 314 (preoccupied—renamed *brunettiana* Alexander, 1920) . W. Himalayas Garhwal District.
- **T. (B.) tessellatipennis* Brun., 1912 : 317 W. Himalayas : Kumaon.
- **T. (Nippotipula) pulcherrima* Brun., 1912 : 310 W. Himalayas : Kumaon.
- **T. (Schummelia) continuata* Brun., 1912 : 328 E. Himalayas : Darjiling.
- **T. (S.) demarcata* (Brun., 1912 : 344, as *Pachyrhina*) (preoccupied—renamed *sessilis* Edwards) E. Himalayas : Darjiling.
- **T. (S.) picticornis* (Brun., 1918 : 279, as *Nesopeza*) Burma : Southern Shan States.
- **T. (Formotipula) rufiventris* Brun., 1918 : 268 (preoccupied—renamed *rufobrachialis* Alexander, 1927) E. Himalayas : Darjiling.
- T. (Yamatotipula) sumifasciata* Brun., 1911 : 250 (synonym of *T. (Y.) nova* Walker, 1848) Assam : Manipur State.
- T. (Acutipula) interrupta* Brun., 1911 : 256 E. Himalayas : Darjiling.

- **T. (A.) munda* Brun., 1912 336 (synonym *T. (A.) vicaria* Walker, 1856, preoccupied) W. Himalayas : Mussoorie.
- **T. (A.) princeps* Brun., 1912 : 306 (synonym *T. (A.) fuscinervis* Brun., 1912 312) E. Himalayas Kurseong.
- **T. (A.) quadrinotata* Brun., 1912 : 330 (synonym *T. (A.) fumicosta* Brun., 1918 266) Assam : Manipur State.
- T. (A.) robusta* Brun., 1911 254 (synonyms) E. Himalayas Darjiling.
- T. (A.) fumifascipennis* Brun., 1918 266 ; W. Himalayas Mussoorie.
 T. (A.) nigrotibialis Brun., 1912 324)
- **T. (A.) fumifascipennis* Brun., 1918 266 E. Himalayas : Darjiling.
- **T. (Indotipula) cinctoterminalis* Brun., 1912 338 E. Himalayas Kurseong.
- **T. (I.) demarcata* Brun., 1911 259 Ceylon.
- T. (I.) divisa* Brun., 1911 261 E. Himalayas Darjiling.
- **T. (I.) elegantula* Brun., 1912 : 339 Assam.
- **T. (I.) flavescens* Brun., 1912 334 Ceylon.
- **T. (I.) flavithorax* Brun., 1918 : 268 S. India : Cochin State.
- T. (I.) gracilis* Brun., 1911 262 E. Himalayas Darjiling.
- **T. (I.) walkeri* Brun., 1911 : 246 Nepal.
- (renaming of *T. (I.) fulvipennis* Walker, 1850, pre-occupied ; synonym **T. (I.) tenuipes* Brun., 1912 333) Assam Sylhet.
- **T. (Tipulodina) contigua* Brun., 1918 265 Assam Garo Hills.
- **T. (T.) gracillima* Brun., 1912 302 Ceylon.
- **T. (T.) patricia* Brun., 1912 : 313 E. Himalayas : Kurseong.
- **T. (T.) simillima* Brun., 1918 265 S. W. India North Kanara District.
- **T. (Vestiplex) brevis* Brun., 1918 : 270 Assam : Khasi Hills.
- (synonym of *T. (V.) reposita* Walker, 1848) Nepal.
- **T. (V.) himalayensis* Brun., 1911 252 . E. Himalayas : Darjiling.
- T. (V.) nigroapicalis* Brun., 1911 257 E. Himalayas : Darjiling.
- **T. (V.) quasimarmoratipennis* Brun., 1912 : 320 E. Himalayas : Darjiling.
- **T. (V.) subtincta* Brun., 1912 : 326 E. Himalayas : Kurseong.
- **T. (Oreomyza) striatipennis* Brun., 1912 : 325 E. Himalayas : Kurseong.
- T. (Lunatipula) fasciculata* Brun., 1918 : 269 pre-occupied—renamed *subernalis* Alexander, 1927) E. Himalayas : Darjiling.
- **T. (L.) marmoratipennis* Brun., 1912 : 319 E. Himalayas : Darjiling.
- **T. (Subgenus incertis) brunnicosta* Brun., 1912 332 W. Himalayas : Simla.

- T. (Subgenus incertis) *elegans* Brun., 1911 : 255
 (preoccupied—renamed *perelegans* Alexander,
 1921) W. Himalayas : Mussoorie.
- *T. (Subgenus incertis) *filicornis* Brun., 1918 : 267 E. Himalayas : Darjiling.
- **Dolichopeza (Oropeza) albatarsis* (Brun., 1918 : 278,
 as *Nesopeza*) E. Himalayas : Darjiling.
- *D. (*Sinoropeza*) *postica* Brun., 1912 : 564 E. Himalayas Darjiling.
- *D. (*Mitopeza*) *longicornis* (Brun., 1918 : 278, as
Nesopeza) Assam Garo Hills.
- *D. (*Nesopeza*) *costalis* Brun., 1918 : 277 S. India Cochin State.
- *D. (*N.*) *infuscata* Brun., 1912 : 565 S. India : Nilgiri Hills.
- D. (*N.*) *obscura* Brun., 1912 : 355
 (preoccupied—renamed *himalayae* Alexander,
 1952) E. Himalayas Kurseong.
- D. (*N.*) *orientalis* Brun., 1912 : 354 E. Himalayas Kurseong.

Nesopeza picticornis Brun., 1918 : 279
 see *Tipula (Schummelia) picticornis* Brun.

CYLINDROTOMINAE

- **Stibadocera quadricellula* (Brun., 1911 : 268, as
Cylindrotoma) E. Himalayas : Kurseong.
- **Stibadocerella pristina* Brun., 1918 : 283 (synonyms
Cylindrotoma latefurcata Brun., 1918 : 280
 lapsus *Stibadocerella bifurcata* Brun., 1920
 212 ; lapsus) Assam Garo Hills.

LIMONIINAE

LECHRIINI

- **Lechria bengalensis* Brun., 1911 : 301 Calcutta ; Pusa.
- **Trichoneura (Xipholimnobia) nepalensis* (Brun.,
 1918 : 317, as *Lechria*) Nepal.

LIMONIINI

- Limonia (Limonia) biannulata* (Brun., 1912 : 430, as
Teucholabis) E. Himalayas : Kurseong.
- *L. (*L.*) *bicolor* (Brun., 1918 : 285, as *Dicranomyia*) Assam : Khasi Hills.
 (synonym *L. (L.) bipunctata* Brun., 1918 : 291,
 as *Limnobia*) Assam : Garo Hills.
- *L. (*L.*) *centralis* (Brun., 1912 : 403, as *Limnobia*) E. Himalayas : Kurseong.
- L. (*L.*) *cinctiventris* (Brun., 1912 : 382, as *Dicranomyia*) E. Himalayas : Kurseong.
 (synonym—*L. (L.) vitripennis* Brun., 1912 : 405,
 as *Limnobia*) E. Himalayas : Darjiling.
- *L. (*L.*) *confinis* (Brun., 1918 : 290, as *Limnobia*) Assam : Garo Hills.
 (preoccupied—renamed *brunettiella* Alexander,
 1929)
- *L. (*L.*) *festiva* (Brun., 1912 : 400, as *Limnobia*) E. Himalayas : Kurseong.

- **L. (L.) flavocincta* (Brun., 1918 : 289, as *Limnobia*) . W. India : Satara Distr.
- **L. (L.) holoptica* (Brun., 1911 : 273, as *Atypoph-thalmus*) Calcutta.
(synonym of *L. (L.) umbrata* de Meijere, 1911; synonym *L. (L.) quinque-notata* Brun., 1918 : 292, as *Limnobia*) Assam : Garo Hills.
- **L. (L.) longipennis* (Brun., 1918 : 292, as *Limnobia*)
(preoccupied—renamed *garoensis* Alexander, 1921) Assam : Garo Hills.
- **L. (L.) marginata* (Brun., 1918 : 290, as *Limnobia*)
(preoccupied—renamed *marginella* Alexander, 1929) Assam : Garo Hills.
- **L. (L.) nigra* (Brun., 1912 : 404, as *Limnobia*)
(preoccupied—renamed *brunettii* Alexander, 1921). May prove to be identical with *bidentata* (Skuse, 1890) S- India : Travancore.
- **L. (L.) nigrescens* (Brun., 1918 : 293, as *Limnobia*)
(preoccupied—renamed *nigricans* Alexander, 1929) Assam : Garo Hills.
- **L. (L.) niveipes* (Brun., 1912 : 404, as *Limnobia*) E. Himalayas : Darjiling.
- **L. Libnotes fuscinervis* (Brun., 1912 : 411, as *Libnotes*) E. Himalayas : Darjiling.
- **L. (L.) indica* (Brun., 1912 : 401, as *Limnobia*) Calcutta.
- **L. (L.) longinervis* (Brun., 1912 : 403, as *Limnobia*) E. Himalayas : Kurseong.
- **L. (L.) notatinervis* (Brun., 1912 : 412, as *Libnotes*) E. Himalayas : Kurseong.
- **L. (L.) punctithorax* (Brun., 1918 : 293, as *Limnobia*) SW. India : N. Kanara District
- **L. (L.) trimaculata* (Brun., 1912 : 402, as *Limnobia*) E. Himalayas : Kurseong.
- **L. (L.) tritincta* (Brun., 1918 : 291, as *Limnobia*) Assam : Garo Hills.
- **L. (Dicranomyia) absens* (Brun., 1912 : 372, as *Dicranomyia*) E. Himalayas : Kurseong.
- **L. (D.) approximata* (Brun., 1912 : 567, as *Dicranomyia*) E. Himalayas : Darjiling.
- **L. (D.) cinerascens* (Brun., 1912 : 381, as *Dicranomyia*) E. Himalayas : Darjiling.
- **L. (D.) columbina* (Brun., 1912 : 567, as *Dicranomyia*) Ceylon.
- **L. (D.) delicata* (Brun., 1912 : 383, as *Dicranomyia*) E. Himalayas : Darjiling.
- **L. (D.) fascipennis* (Brun., 1912 : 379, as *Dicranomyia*) E. Himalayas : Kurseong.
- **L. (D.) flavobrunnea* (Brun., 1912 : 384, as *Dicranomyia*) Calcutta.
- **L. (D.) fortis* (Brun., 1912 : 385, as *Dicranomyia*) E. Himalayas : Darjiling.
- **L. (D.) fraterna* (Brun., 1912 : 378, as *Dicranomyia*) E. Himalayas : Darjiling.

- **L. (D.) nigrithorax* (Brun., 1912 : 385, as *Dicranomyia*) E. Himalayas : Darjiling.
- **L. (D.) pulchripennis* (Brun., 1912 : 376, as *Dicranomyia*) E. Himalayas . Darjiling.
- **L. (D.) puncticosta* (Brun., 1912 : 377, as *Dicranomyia*) E. Himalayas : Kurseong.
- (synonym *L. (L.) tinctinervis* Brun., 1912 : 401, as *Limnobia*) E. Himalayas: Darjiling.
- **L. (D.) simplex* (Brun., 1912 : 384, as *Dicranomyia*) Calcutta : Assam.
- **L. (D.) sordida* (Brun., 1912 : 382, as *Dicranomyia*) E. Himalayas : Kurseong.
- (synonym *L. (L.) innocens* Brun., 1912 : 568, as *Dicranomyia*) W. Himalayas : Almora, Kumaon.
- **L. (D.) subfascipennis* (Brun., 1912 : 380, as *Dicranomyia*) E. Himalayas : Kurseong.
- **L. (Geranomyia) circipunctata* (Brun. 1912 : 390, as *Geranomyia*) India : Lower Bengal.
- **L. (G.) flavigosta* (Brun., 1912 : 389 as *Geranomyia*) India : Ganges Delta.
- **L. (G.) flaviventris* (Brun., 1918 : 288, as *Geranomyia*) E. Himalayas: Darjiling.
- L. (G.) genitalis* (Brun., 1911 : 275, as *Geranomyia*) S. India: Tenmalai, Travancore.
- (synonym of *L. (G.) fletcheri* Edwards, 1911)
- L. (G.) nigronotata* (Brun., 1918 : 287, as *Geranomyia*) S. India : Malabar District.
- **L. (G.) notatipennis* (Brun., 1913a : 152, as *Geranomyia*) Assam Abor District.
- **L. (G.) pulchripennis* (Brun., 1912 : 393, as *Geranomyia*) E. Himalayas : Kurseong.
- (preoccupied—renamed *pictorum* Alexander, 1929)
- L. (G.) semifasciata* (Brun., 1911 : 276, as *Geranomyia*) E. Himalayas : Darjiling.
- **L. (G.) semiannulata* (Brun., 1911 : 277, as *Geranomyia*) Bihar : Paresnath.
- **L. (G.) tridens* (Brun., 1912 : 391, as *Geranomyia*) India : Ganges Delta.
- L. (G.) vinaceobrunnea* (Brun., 1911 : 274, as *Geranomyia*) W. Himalayas : Simla.
- **L. (Rhipidia) antennata* (Brun., 1911 : 272, as *Ceratostephanus*) W. Himalayas : Simla.
- **L. (R.) demarcata* (Brun., 1912 : 370, as *Dicranomyia*) E. Himalayas : Kurseong.

- **L. (R.) marmoripennis* (Brun., 1912 : 369, as *Dicranomyia*) E. Himalayas : Kurseong.
 (synonym of *L. (R.) pulchra* de Meijere)
- **L. (R.) subtesselata* (Brun., 1912 : 565, as *Dicranomyia*) (synonym *L. (R.) zeylanica* S. W.) . . Ceylon.
- **L. (Alexandriaria) prominens* (Brun., 1918 : 285, as *Dicranomyia*) . . . SW. India : Goa.
- **L. (Pseudoglochina) bicinctipes* (Brun., 1912 : 566, as *Dicranomyia*) . . . Burma : Dawna Hills.
- **L. (P.) pictipes* (Brun., 1918 : 286, as *Dicranomyia*) SW. India : Goa.
Dicranomyia cinctiventris Brun., 1912 : 382 . . . see *Limonia (Limonia)*.
- D. demarcata* Brun., 1912 : 370 . . . see *Limonia (Rhipidia)*.
- **D. festiva* Brun., 1912 : 400 . . . see *Limonia (Limonia)*.
- D. marmoripennis* Brun., 1912 : 369 . . . see *Limonia (Rhipidia)*.
- **D. niveiapicalis* Brun., 1918 : 285 . . . see *Riedelomyia*.
- D. ornatipes* Brun., 1912 : 380 see *Gonomyia (Lipophleps)*.
- D. prominens* Brun., 1918 : 285 see *Limonia (Alexandriaria)*.
- **Limnobia triangularis* Brun., 1912 : 406 . . . see *Antocha (Antocha)*.
- L. trimaculata* Brun., 1912 : 402 see *Limonia (Libnotes)*.
- **Helius (Helius) ferruginosus* (Brun., 1912 : 418, as *Rhamphidia*) Burma : Dawna Hills.
- H. (H.) fratellus* (Brun., 1918 : 296, as *Rhamphidia*) S.W. India : N. Kanara District.
- **H. (H.) inconspicuus* (Brun., 1912 : 419, as *Rhamphidia*) E. Himalayas : Kurseong.
- **H. (H.) unicolor* (Brun., 1912 : 419, as *Rhamphidia*) E. Himalayas : Darjiling.
- **H. (Eurhamphidia) abnormalis* (Brun., 1918 : 296, as *Rhamphidia*) Assam : Khasi Hills.
- **Antocha (Antocha) indica* Brun., 1912 : 426 . . . E. Himalayas : Kurseong.
- **A. (A.) triangularis* (Brun., 1912 : 406, as *Limnobia*) W. Himalayas : Simla.
- **A. (A.) unilineata* Brun., 1912 : 427 W. Himalayas : Dehra Dun.
- **Orimarga (Orimarga) peregrina* Brun., 1912 : 424 E. Himalayas : Darjiling.

PEDICINNI

- **Pedicia (Tricyphona) elegans* (Brun., 1912 : 516, as *Amalopis*) E. Himalayas : Kurseong.
- P. (T.) flavigennnis* (Brun., 1918 : 327, as *Limnophila*) . . E. Himalayas : Darjiling.
- **P. (T.) glabripennis* (Brun., 1912 : 515, as *Amalopis*) E. Himalayas : Darjiling.

- * *Dicranota (Amalopina) elegantula* (Brun., 1912 : 517,
as *Amalopina*) E. Himalayas : Darjiling.
- D. (Plectromyia) incompleta* (Brun., 1912 : 493, as
Gnophomyia) E. Himalayas : Kurseong.
- * *D. (Raphidolabina) spectralis* (Brun., 1918 : 321,
as *Amalopis*) E. Himalayas : Darjiling.
- * *D. (Raphidolabis) aperta* (Brun., 1912 : 492, as
Gnophomyia) E. Himalayas : Darjiling.
- (preoccupied—renamed *brunettii* Edwards,
1916)
- * *D. (R.) fascipennis* (Brun., 1911 : 289, as *Claduroides*) E. Himalayas : Darjiling.
- (synonyms *D. (R.) fascipennis* Brun., 1912 :
519, as *Raphidolabis* ; *D. (R.) longipennis*
(Brun., 1912 : 489, as *Gnophomyia*)
- D. (R.) sordida* (Brun., 1911 : 290, as *Claduroides*)
(synonym **D. (R.) indica* (Brun., 1912 : 519,
as *Raphidolabis*) W. Himalayas : Simla.

HEXATOMINI

- * *Paradelphomyia (Oxyrhiza) flavescentis* (Brun., 1911 :
284, as *Cladura*) E. Himalayas : Darjiling.
- (synonym—*P. (O.) furcata* (Brun., 1912 : 491,
as *Gnophomyia*)
- * *Epiphragma (Epiphragma) kempfi* Brun., 1913a :
155 NE. Indian Frontier.
- * *E. (E.) ornatipennis* (Brun., 1918 : 328, as *Limno-*
phila) E. Himalayas : Darjiling.
- * *E. (E.) vicina* Brun., 1918 : 331 E. Himalayas : Darjiling.
- * *Taiwanomyia cavernicola* (Brun., 1924 : 100, as
Troglophila) Assam : Siju Cave, Garo Hills.
- Pseudolimnophila (Pseudolimnophila) fusca* (Brun.,
1918 : 326, as *Limnophila*) E. Himalayas : Darjiling.
- * *P. (P.) glabra* (Brun., 1918 : 325, as *Limnophila*).
May belong in *Pilaria* Sintenis SW. India : N. Kanara District.
- * *P. (P.) multipunctata* (Brun., 1912 : 569 : as *Limno-*
phila) S. India : Nilgiri Hills.
- P. (P.) pallidicoxa* (Brun., 1912 : 523, as *Limnophila*) E. Himalayas : Darjiling.
- * *Eupilaria annulipes* (Brun., 1918 : 324, as *Limnophila*) SW. India : N. Kanara District.
- * *E. inconsequens* (Brun., 1918 : 326, as *Limnophila*) SW. India : N. Kanara District.
- Limnophila (Elaeophila) fascipennis* (Brun., 1912 :
526, as *Ephelia*) E. Himalayas : Darjiling.
- L. (E.) ornata* (Brun., 1912 : 527, as *Ephelia*) W. Himalayas : Kumaon.

- **L. (Dicranophragma) interrupta* (Brun., 1918 : 319,
as *Cladura*) E. Himalayas : Darjiling.
- L. (D.) multipunctipennis* (Brun., 1918 : 329, as
Dicranophragma) E. Himalayas : Darjiling.
- **L. (D.) pulchripennis* (Brun., 1912 : 524, as *Dicra-*
nophragma) E. Himalayas : Kurseong.
(preoccupied—renamed *venustipennis* Alexan-
der, 1921)
- **L. (Subgenus incertis) honesta* Brun., 1912 : 570 W. Himalayas : Kumaon.
- **Limnophila annulipes* Brun., 1918 : 324 see *Eupilaria*.
- **L. claripennis* Brun., 1913a : 153 see *Crypteria*.
- L. flavigennis* Brun., 1918 : 327 see *Pedicia (Tricyphona)*.
- **L. fusca* Brun., 1918 : 326 see *Pseudolimnophila*.
- **L. glabra* Brun., 1918 : 325 see *Pseudolimnophila*.
- **L. incompleta* Brun., 1918 : 326 see *Gonomyia (Protogonomyia)*.
- **L. inconsequens* Brun., 1918 : 326 see *Eupilaria*.
- **L. multipunctata* Brun., 1912 : 569 see *Pseudolimnophila*.
- **L. ornatipennis* Brun., 1918 : 328 see *Epiphragma*.
- L. pallidicoxa* Brun., 1912 : 523 see *Pseudolimnophila*.
- **L. parvicellula* Brun., 1918 : 325 see *Gonomyia (Gonomyia)*.
- **L. quartaria* Brun., 1913a : 154 see *Gnophomyia*.
- **L. simplex* Brun., 1912 : 523 see *Neolimnophila*.
- **Hexatoma (Eriocera) aterrima* (Brun., 1912 : 540,
as *Eriocera*) S. India : Travancore.
- **H. (E.) aurantia* (Brun., 1918 : 335, as *Eriocera*) E. Himalayas : Darjiling.
- H. (E.) badia* (Brun., 1911 : 310, as *Eriocera*) . Ceylon.
- **H. (E.) caliginosa* (Brun., 1918 : 339, as *Eriocera*) S. India : N. Kanara District.
- H. (E.) cincta* (Brun., 1918 : 333, as *Eriocera*) E. Himalayas : Darjiling District.
(synonym **H. (E.) cingulata* Brun., 1912 :
570, preoccupied)
- **H. (E.) decorata* (Brun., 1918 : 337, as *Eriocera*) . E. Himalayas : Darjiling.
- **H. (E.) elongatissima* (Brun., 1912 : 542, as
Eriocera) S. India : Travancore.
- H. (E.) fenestrata* (Brun., 1911 : 312, as *Eriocera*) . Central Tonkin (type) ; Assam :
Khasi Hills.
- **H. (E.) flavipes* (Brun., 1912 : 544, as *Eriocera*) . E. Himalayas : Kurseong.
- **H. (E.) gravelyi* (Brun., 1918 : 337, as *Eriocera*) . E. Himalayas : Darjiling.

- **H. (E.) greenii* (Brun., 1911 : 313, as *Eriocera*) . . . Ceylon.
- **H. (E.) kempfi* (Brun., 1918 : 339, as *Eriocera*) . . . SW. India : N. Kanara District.
- **H. (E.) nigerrima* (Brun., 1912 : 571, as *Eriocera*) . . . E. Himalayas : Darjiling, Ghumti.
- **H. (E.) plumbicincta* (Brun., 1911 : 311, as *Eriocera*) . . . Assam : Manipur.
- **H. (E.) pulchrithorax* (Brun., 1918 : 336, as *Eriocera*) . . . S. India : Cchin State.
- H. (E.) rufibasis* (Brun., 1911 : 310, as *Eriocera*) . . . Lower Burma : Tenasserim.
- **H. (E.) rufithorax* (Brun., 1912 : 534, as *Eriocera*) . . . Ceylon.
(Synonym of *H. (E.) ctenophoroides* Edwards, 1911)
- **H. (E.) rufiventris* (Brun., 1918 : 336, as *Eriocera*) . . . S. India : Coorg ; Cochin State.
- H. (E.) semilimpida* (Brun., 1911 : 311, as *Eriocera*) . . . Assam : Khasi Hills.
(synonym **H. (E.) maculiventris* Brun., 1918 : 340, as *Eriocera*) . . . Assam : Garo Hills.
- **H. (E.) tenuis* (Brun., 1912 : 539, as *Eriocera*) . . . S. India : Nilgiri Hills.
- **H. (E.) testacea* (Brun., 1912 : 548, as *Eriocera*).
May be identical with *tenuis* . . . S. India : Nilgiri Hills.
- **H. (E.) triangularis* (Brun., 1912 : 572, as *Eriocera*) . . . S. India : Nilgiri Hills.
- **H. (E.) tripunctipennis* (Brun., 1918 : 338, as *Eriocera*) . . . SW. India : N. Kanara District.
- **Eriocera nigroapicalis* Brun., 1918 : 335 . . . see *Nephrotoma*.
- **Atarba (Atarbodes) flava* Brun., 1912 : 435 . . . E. Himalayas : Darjiling.
- **Elephantomyia (Elephantomyodes) aurantia* (Brun., 1918 : 289, as *Aporosa*) . . . Assam : Garo Hills.

ERIOPTERINI

- **Franckomyia gracilis* (Brun., 1913a : 156, as *Dicranophragma*) . . . NE. Indian Frontier.
(preoccupied—renamed *recessiva* Alexander, 1927)
- **Crypteria claripennis* (Brun., 1913a : 153, as *Limnophila*) . . . NE. Indian Frontier.
- Neolimnophila genitalis* (Brun., 1912 : 490, as *Gnophomyia*) (synonym—*N. simplex* Brun., 1912 : 523) . . . W. Himalayas : Kumaon, Bhowali.
- **Dasymallomyia signata* Brun., 1911 : 304 . . . E. Himalayas : Kurseong.
- **Gnophomyia quartaria* (Brun., 1913a : 154, as *Limnophila*) . . . NE. Indian Frontier.
- G. strenua* Brun., 1912 : 492 . . . E. Himalayas : Kurseong.
- **Gnophomyia aperta* Brun., 1912 : 492 . . . see *Dicranota (Rhaphidolabis)*

- G. furcata* Brun., 1912 : 491 see *Paradelphomyia*.
- G. genitalis* Brun., 1912 : 490 see *Neolimnophilla*.
- G. incompleta* Brun., 1912 : 493 see *Dicranota (Plectromyia)*.
- G. longipennis* Brun., 1912 : 489 see *Dicranota (Raphidolabis)*.
- G. nigra* Brun., 1912 : 494 see *Gonomyia (Protagonomyia)*.
- Gonomyia (Protagonomyia) nigripes* (Brun., 1912 : 458, as *Mesocyphona*) E. Himalayas : Darjiling.
- (synonyms *G. (P.) gracilis* (Brun., 1918 : 310, as *Mesocyphona*) E. Himalayas : Assam-Bhutan Frontier.
- **G. (P.) incompleta* (Brun., 1918 : 326, as *Limnophila*) E. Himalayas : Darjiling.
- **G. (P.) nigra* (Brun., 1912 : 494, as *Gnophomyia*) W. Himalayas : Kumaon.
- **G. (Idiocera) proxima* Brun., 1912 : 474 Bengal.
- **G. (Gonomyia) affinis* Brun., 1912 : 472 E. Himalayas : Kurseong.
- **G. (G.) aperta* Brun., 1912 : 473 Bihar : Purneah District.
- **G. (G.) parvicellula* (Brun., 1918 : 325, as *Limnophila*) SW. India : N. Kanara District.
- **G. (Lipophleps) incompleta* Brun., 1912 : 471 Calcutta.
- **G. (L.) flavomarginata* Brun., 1912 : 472 E. Himalayas : Kurseong.
- G. (L.) ornatipes* (Brun., 1912 : 380, as *Dicranomyia*) S. India : Travancore.
- **Gonomyia antica* Brun., 1912 : 568 see *Cheilotrichia (Empeda)*.
- **Teucholabis (Teucholabis) angusticapitis* Brun., 1918 : 305 Assam : Garo Hills.
- **T. (T.) assamensis* Brun., 1918 : 306 Assam : Khasi Hills.
- **T. (T.) fenestrata insignis* Brun., 1912 : 430 S. India : Travancore.
- **T. (T.) ornata* Brun., 1918 : 305 Ceylon.
- **Teucholabis biannulata* Brun., 1912 : 430 see *Limonia (Limonia)*.
- **Gymnastes (Gymnastes) violacea* Brun., 1911 : 282
(synonym of *G. (G.) cyanea* Edwards, 1911 : 61) Ceylon.
- **G. (Paragymnastes) bistriatipennis* Brun., 1918 : 307 SW. India : N. Kanara District.
- G. (P.) pennipes* Brun., 1918 : 308 Assam : Garo Hills.
- **Trentepohlia (Plesiomongoma) venosa* (Brun., 1918 : 314, as *Plesiomongoma*) Assam : Khasi Hills.
- **T. (Mongoma) flava* (Brun., 1918 : 314, as *Mongoma*) Assam : Garo Hills.
- **T. (M.) kempfi* (Brun., 1918 : 313, as *Mongoma*) Assam : Garo Hills.

- **T. (M.) pallidiventris* (Brun., 1912 : 481, as *Mongoma*) S. India : Travancore.
(may be identical with *T. (M.) tenera* O.S., 1882 : 89)
- **T. (M.) splendida* (Brun., 1918 : 313, as *Mongoma*) Assam : Garo Hills.
- T. (Anchimongoma) simplex* (Brun., 1918 : 316, as
Anchimongoma) S. India : N. Kanara District.
- **T. (Trentepohlia) albogeniculata* (Brun., 1912 : 569,
as *Mongomiooides*) Burma : Dawna Hills.
- **T. (T.) marmorata* (Brun., 1912 : 483, as *Mongo-*
miooides) Calcutta.
- **T. (T.) nigroapicalis* (Brun., 1912 : 483, as *Mongo-*
miooides) Lucknow.
- **T. (T.) ornatipennis* Brun., 1918 : 315 SW. India : N. Kanara District.
- **Riedelomyia niveapicalis* (Brun., 1918 : 285, as
Dicranomyia) SW. India : N. Kanara District.
- **Ormosia (Ormosia) geniculata* (Brun., 1912 : 441,
as *Rhynopholopha*) E. Himalayas : Kurseong.
- **O. (O.) pulchra* (Brun., 1912 : 442, as *Rhynopholopha*) W. Himalayas : Phagu, Simla
District.
- **Cryptolabis (Baeoura) distans* (Brun., 1912 : 451,
as *Erioptera*) E. Himalayas : Kurseong.
- Cheilotrichia (Empeda) brevior* (Brun., 1912 : 452, E. Himalayas : Darjiling.
as *Erioptera*)
- (synonyms: *C. (E.) antica* (Brun., 1912 : 568,
as *Gonomyia*) E. Himalayas : Ghumti.
- C. (E.) inconspicua* (Brun., 1912 : 475, as
Empeda) E. Himalayas : Darjiling.
- **Erioptera (Meteriopelta) flava* Brun., 1912 : 455 Bengal : Calcutta.
(preoccupied—renamed *bengalensis* Alexander,
1921)
- **E. (M.) halterata* Brun., 1912 : 457 E. Himalayas : Darjiling.
- **E. (M.) subincta* Brun., 1912 : 455 E. Himalayas : Darjiling.
- **E. (Erloptera) ferruginea* Brun., 1912 : 450 S. India : Travancore State.
- **E. (E.) grandior* Brun., 1912 : 456 W. Himalayas : Simla.
- **E. (E.) incerta* Brun., 1912 : 452 E. Himalayas : Darjiling.
- **E. (E.) orientalis* Brun., 1912 : 453 (synonym *E.*
(E.) dictenidia Alexander, 1921) E. Himalayas : Darjiling.
- **E. (Teleneura) parallela* Brun., 1912 : 453 E. Himalayas : Kurseong.
- **E. (T.) punctipennis* Brun., 1912 : 449 E. Himalayas : Kurseong.
(preoccupied—renamed *annandaleana* Alexander,
1953)

- **Erioptera genitalis* Brun., 1912 : 456 . . . see *Molophilus*.
- **Molophilus (Molophilus) assamensis* Brun., 1912 : 445 . . . Assam : Sylhet.
- **M. (M.) genitalis* (Brun., 1912 : 456, as *Erioptera*) . W. Himalayas : Kumaon, Bhowali.
- **M. (M.) inconspicuus* Brun., 1912 : 444 . W. Himalayas : Simla.
- **Styringomyia flava* Brun., 1911 : 301 . S. India : Travancore State.
- **S. obscura* Brun., 1911 : 300 . Nepal : Thamaspur.
- **Toxorhina (Ceratocheilus) brevifrons* (Brun., 1918 : 300, as *Conifrons*) . Assam : Garo Hills.
- T. (Toxorhina) incerta* Brun., 1912 : 422 . E. Himalayas : Kurseong.

III—THE EXTRA-LIMITAL TIPULIDAE DESCRIBED BY BRUNETTI

TIPULINAE

Ctenacroscelis borneensis (Brun., 1918 : 263, as *Tipula*) Borneo.
(renaming of *Tipula pallida* Walker, 1865, preoccupied)

Ctenophora (Pselliophora) bifascipennis (Brun., 1911 : 241, as *Pselliophora*) E. China : Shanghai.

(synonyms)—

C. (P.) sackeni Edwards, 1916 : 257 . as *Pselliophora*.

C. (P.) horikawae Matsumura, 1916 : 449 as *Psehicphora*.

C. (P.) compta Enderlein, 1921 : 220 . as *Pselliophora*.

C. (Pselliophora) terminalis (Brun., 1911 : 244, as *Pselliophora*) Indochina : Tonkin.

Tipula (Formotipula) sciariformis Brun., 1911 : 259 . Indochina : Tonkin.

T. (Tipulodina) schummeli Brun., 1911 : 247 . Amboina.
(synonym of *T. (T.) sumifinis* Walker, 1861 : 145, synonym—*T. (T.) longicornis* Doleschall, 1858 : 79, preoccupied)

T. (Subgenus incertis) cinerea Brun., 1911 : 252 Lombok.
(preoccupied—renamed *lombokensis* Alexander, 1942)

T. (Subgenus incertis) pluto Brun., 1911 : 251 . . . Indochina : Tonkin.

LIMONIINAE

**Orimarga (Orimarga) borneensis* Brun., 1911 : 280 . Borneo : Sarawak.

Gynoplistia (Gynoplistia) octofasciata Brun., 1911 : 307 South Celebes.

**Toxorhina (Ceratocheilus) latifrons* (Brun., 1918 : 299, *Conithorax*) Malay States : Selangor-Pahang Border.

IV.—SYNOPSIS

The accompanying report discusses the Tipuloidean flies that were described by the late Enrico Brunetti between 1911 and 1924, a total of approximately 240 valid species, with numerous additional names that are considered to be synonyms of other species. This synonymy and the erroneous assignment to genera of many of his species has made it difficult to proceed with studies on the Indian crane-fly fauna. The late Dr. Fred W. Edwards was able to settle much of the synonymy through a study of the personal collection made by Brunetti and now preserved in the British Museum of Natural History. The great bulk of Brunetti's types are preserved in the collections of the Zoological Survey of India at Calcutta and several doubtful species were made available to the present writer for appraisal. As a result of these examinations of critical materials it is believed that the synonymy of the Brunetti species is more firmly established.

The apparently valid species described by Brunetti are distributed in the following groups :

Ptychopteridae	3
Trichoceridae	4
Tipulidae : Tipulinae	71
Cylindrotominae	2
Limoniinae—										
Limoniini	59
Lechriini	2
Pediciini	9
Hexatomini	41
Eriopterini	51
Total										242

Dr. A. P. Kapur has indicated the various types that are preserved in the collection of the Zoological Survey of India.